

2012

Evaluation of Survey Indices for Scaled Quail in West Texas

Barrett A. Koennecke

Rolling Plains Quail Research Ranch

Dale Rollins

Texas A&M University

Chris Snow

Angelo State University

Jeff White

University Lands

Follow this and additional works at: <https://trace.tennessee.edu/nqsp>

Recommended Citation

Koennecke, Barrett A.; Rollins, Dale; Snow, Chris; and White, Jeff (2012) "Evaluation of Survey Indices for Scaled Quail in West Texas," *National Quail Symposium Proceedings*: Vol. 7 , Article 129.

<https://doi.org/10.7290/nqsp07jueq>

Available at: <https://trace.tennessee.edu/nqsp/vol7/iss1/129>

This article is brought to you freely and openly by Volunteer, Open-access, Library-hosted Journals (VOL Journals), published in partnership with The University of Tennessee (UT) University Libraries. This article has been accepted for inclusion in National Quail Symposium Proceedings by an authorized editor. For more information, please visit <https://trace.tennessee.edu/nqsp>.

EVALUATION OF SURVEY INDICES FOR SCALED QUAIL IN WEST TEXAS

Barrett A. Koennecke¹

Rolling Plains Quail Research Ranch, 1262 U.S. Highway 180 West, Rotan, TX 79546, USA

Dale Rollins

Texas AgriLife Research, Texas A&M University System, 7887 U.S. Highway 87 North, San Angelo, TX 76901, USA

Chris Snow

Department of Biology, Angelo State University, 2601 West Avenue N, San Angelo, TX 76909, USA

Jeff White

University Lands, 704 West Dengar Avenue, Midland, TX 79705, USA

ABSTRACT

Wildlife biologists and land managers require information on population demographics to effectively plan harvest schedules and evaluate habitat modifications. Population indices can potentially provide an efficient way to gather reliable information on wildlife populations as long as they reflect population behavior. We evaluated the relationships among standard survey indices used to monitor scaled quail (*Callipepla squamata*) across 6 sites in west Texas from 2007 to 2010. We collected data on spring cock call counts, simulated nest survival, roadside counts, and helicopter counts. The mean difference between methods was 2.4 quail/1.6 km on the Andrews County sites, and 4 quail/1.6 km at the Upton/Reagan County sites. Roadside counts and helicopter counts had similar numerical trends in relative abundance with a correlation coefficient of (0.67). Simulated nest fate (i.e., dummy nests) tended to track trends in population abundance. Our survey indices also followed annual fluctuations in scaled quail abundance as estimated from Texas Parks and Wildlife Department's annual roadside surveys.

Citation: Koennecke, B. A., D. Rollins, C. Snow, and J. White. 2012. Evaluation of survey indices for scaled quail in west Texas. Proceedings of the National Quail Symposium 7:364.

¹E-mail: bkoennecke@ag.tamu.edu