

2012

The Forgotten Quail Decline: The Plight of Scaled Quail in Texas

Fidel Hernandez
Texas A&M University

Chad J. Parent
Texas A&M University

Ian C. Trewella
Texas A&M University

Eric D. Grahmann
Texas A&M University

Follow this and additional works at: <https://trace.tennessee.edu/nqsp>

Recommended Citation

Hernandez, Fidel; Parent, Chad J.; Trewella, Ian C.; and Grahmann, Eric D. (2012) "The Forgotten Quail Decline: The Plight of Scaled Quail in Texas," *National Quail Symposium Proceedings*: Vol. 7 , Article 130.
<https://doi.org/10.7290/nqsp07h255>
Available at: <https://trace.tennessee.edu/nqsp/vol7/iss1/130>

This article is brought to you freely and openly by Volunteer, Open-access, Library-hosted Journals (VOL Journals), published in partnership with The University of Tennessee (UT) University Libraries. This article has been accepted for inclusion in National Quail Symposium Proceedings by an authorized editor. For more information, please visit <https://trace.tennessee.edu/nqsp>.

THE FORGOTTEN QUAIL DECLINE: THE PLIGHT OF SCALED QUAIL IN TEXAS

Fidel Hernández¹

Caesar Kleberg Wildlife Research Institute, Texas A&M University-Kingsville, MSC 218, 700 University Boulevard, Kingsville, TX 78363, USA

Chad J. Parent

Caesar Kleberg Wildlife Research Institute, Texas A&M University-Kingsville, MSC 218, 700 University Boulevard, Kingsville, TX 78363, USA

Ian C. Trewella

Caesar Kleberg Wildlife Research Institute, Texas A&M University-Kingsville, MSC 218, 700 University Boulevard, Kingsville, TX 78363, USA

Eric D. Grahmann

Caesar Kleberg Wildlife Research Institute, Texas A&M University-Kingsville, MSC 218, 700 University Boulevard, Kingsville, TX 78363, USA

ABSTRACT

Several quail species are experiencing range-wide declines in the United States. The northern bobwhite (*Colinus virginianus*) has garnered the most attention, both from a research and conservation perspective. The bobwhite decline in Texas, has resulted in considerable time and effort being devoted to research and management on the species due to its status as a highly popular and economically important game bird. This attention has been beneficial to bobwhite conservation and management but, an unfortunate consequence of this focus has been neglect of the scaled quail (*Callipepla squamata*) that has been declining at even a more alarming rate. Scaled quail, according to Breeding Bird Survey (BBS) data, have declined at a rate of 5.1% per year in the Tamaulipan Biotic Province (southern Texas), the greatest of any region surveyed in its geographic range. Anecdotal reports of landowners have long noted the gradual disappearance of scaled quail and concomitant replacement with northern bobwhite throughout southern Texas, beginning since about the 1990s. Analysis of BBS data provides evidence for this replacement. Percent of quail detections in the core of the scaled quail range in the Tamaulipan Biotic Province were 80:20 (scaled quail: bobwhite) during the 1960s but currently represent about 5:95. In addition, the range of scaled quail has been contracting, moving progressively west with time. The species is no longer detected on the easternmost BBS routes in southern Texas.

Citation: Hernández, F., C. J. Parent, I. C. Trewella, and E. D. Grahmann. 2012. The forgotten quail decline: the plight of scaled quail in Texas. Proceedings of the National Quail Symposium 7:365.

Key words: Breeding Bird Survey data, *Callipepla squamata*, *Colinus virginianus*, northern bobwhite, scaled quail

¹E-mail: fidel.hernandez@tamuk.edu