

2017

Genetic Structure of Northern Bobwhite in the Rolling Plains

Peter E. Schlichting
Texas Tech University

Melanie Murphy
University of Wyoming

Byron R. Buckley
Texas Tech University

C. Brad Dabbert
Texas Tech University

Follow this and additional works at: <https://trace.tennessee.edu/nqsp>



Part of the [Natural Resources and Conservation Commons](#)

Recommended Citation

Schlichting, Peter E.; Murphy, Melanie; Buckley, Byron R.; and Dabbert, C. Brad (2017) "Genetic Structure of Northern Bobwhite in the Rolling Plains," *National Quail Symposium Proceedings*: Vol. 8 , Article 83.

<https://doi.org/10.7290/nqsp08wi8m>

Available at: <https://trace.tennessee.edu/nqsp/vol8/iss1/83>

This article is brought to you freely and openly by Volunteer, Open-access, Library-hosted Journals (VOL Journals), published in partnership with The University of Tennessee (UT) University Libraries. This article has been accepted for inclusion in National Quail Symposium Proceedings by an authorized editor. For more information, please visit <https://trace.tennessee.edu/nqsp>.

GENETIC STRUCTURE OF NORTHERN BOBWHITE IN THE ROLLING PLAINS

Peter E. Schlichting^{1,2}

Department of Natural Resources Management, Texas Tech University, Goddard Building, Box 42125, Lubbock, TX 79409, USA

Melanie Murphy

Department of Ecosystem Science and Management, University of Wyoming, Agriculture Building 2013, 1000 E. University Ave. Laramie, WY 82071, USA

Byron Buckley

Department of Natural Resources Management, Texas Tech University, Goddard Building, Box 42125, Lubbock, TX 79409, USA

C. Brad Dabbert

Department of Natural Resources Management, Texas Tech University, Goddard Building, Box 42125, Lubbock, TX 79409, USA

ABSTRACT

The recent declines in northern bobwhite quail populations in the Rolling Plains of Texas have raised concerns about habitat connectivity and gene flow. In addition, bobwhites have several life history traits that make them likely to display high levels of spatial genetic structure including low survival, high reproductive rates, and low dispersal rates. To determine if populations within the Rolling Plains have limited gene flow, we investigated the genetic structure of northern bobwhites within the ecoregion. Blood samples were collected at 16 ranches, encompassing 22 million acres, between February 2010 and April 2013. Bobwhites were also sampled at a ranch in South Texas to serve as an outgroup. Samples ($n = 647$) were genotyped at 14 microsatellite loci that averaged 19.00 ± 5.07 alleles per loci. Global F_{ST} indicated significant genetic structure ($p = 0.001$) between ranches with no isolation by distance signal ($p = 0.079$). Program STRUCTURE, however, indicated many ($n = 30$) overlapping subpopulations with no ranch constituting a single subpopulation and individuals from the outgroup ranch were included in 11 subpopulations. It appears that bobwhites within the Rolling Plains have few restrictions to gene flow and dispersal is not limited by the dominant habitat, xeric rangeland. These results suggest that populations in the Rolling Plains are not in danger of becoming isolated nor are bottlenecks present due to the recent decline.

Citation: Schlichting, P.E., M. Murphy, B. Buckley, and C. B. Dabbert. 2017. Genetic Structure of northern bobwhite in the Rolling Plains. National Quail Symposium Proceedings 8:310.

Key words: Gene flow, connectivity, population decline, Rolling Plains

¹ E-mail: pschlichting@gmail.com

² Present address: Savannah River Ecology Lab, P.O. Drawer E, Aiken, SC 29802

©2017 [Schlichting, Murphy, Buckley and Dabbert] and licensed under CC BY-NC 4.0.