

2009

Index

Follow this and additional works at: <https://trace.tennessee.edu/nqsp>

Recommended Citation

(2009) "Index," *National Quail Symposium Proceedings*: Vol. 6 , Article 55.

Available at: <https://trace.tennessee.edu/nqsp/vol6/iss1/55>

This article is brought to you freely and openly by Volunteer, Open-access, Library-hosted Journals (VOL Journals), published in partnership with The University of Tennessee (UT) University Libraries. This article has been accepted for inclusion in National Quail Symposium Proceedings by an authorized editor. For more information, please visit <https://trace.tennessee.edu/nqsp>.

Index

A

Alectoris rufa, 267
 abundance, 178, 220, 251
 action plan, 291
 active management, 370
 adaptive management, 2
 age, 135
 ageing, 505
 agricultural intensification, 111
 agriculture, 172, 232
 Amdro[®], 78
 Ames Plantation, 121
 area curves, 489
 Arizona, 370
 Arkansas, 146, 184
 Austria, 410
 autumn, 178

B

Bonasa umbellus, 480, 499
 barriers, 322
 basal area, 178
 Belize, 99
 biological, 322
 black-throated bobwhite, 99
 breeding success, 267
 brood, 88
 brood size, 267
 brood survival, 410
 broods, 302
 buffer strips, 232

C

Callipepla gambelii, 106
Callipepla squamata, 210, 456
Colinus nigrogularis, 99
Colinus virginianus, 16, 46, 54, 66, 78, 88, 121, 135, 146, 160, 178, 184, 193, 220, 251, 256, 271, 282, 315, 328, 338, 348, 370, 396, 432, 447, 467, 518
Cyrtonyx montezumae, 322, 426
 cactus, 256
 call counts, 46
 carcass fat, 499
 Chihuahuan Desert, 456
 Chihuahuan desert, 210
 climate trends, 328
 closed-canopy pine, 135
 compositional analysis, 410
 conservation, 111

Conservation Reserve Program, 302
 crippling loss, 420

D

data collection, 370
 decision making, 2
 density, 46
 discriminant function analysis, 505
 dispersal, 135
 distance sampling, 46
 disturbance, 391

E

east Texas, 16, 66
 edge, 184

F

farming, 172
 farmland, 348
 fat condition, 499
 field borders, 232
 fire, 193, 271, 315
 Florida, 88, 193, 271, 282
 food availability, 160
 food habits, 160
 food plots, 16, 66
 food preferences, 160
 food quantity, 146
 forestland, 348
 formal inference-based recursive modeling (FIRM), 302
 French Mediterranean region, 267

G

Gambel's quail, 106
 game crop, 267
 genetic variability, 121
 Georgia, 46, 135, 282, 338, 348
 GIS, 106, 302
 gray partridge, 24, 27, 291, 380, 391
 guzzlers, 210

H

habitat, 106, 135, 172, 193, 251, 302, 410
 habitat enhancement, 391
 habitat loss, 370
 habitat management, 260, 267
 habitat modeling, 220
 habitat quality, 184
 habitat selection, 88

- habitat use, 232, 391
- half-cutting, 251
- harvest, 260, 420
- herbicide, 315
- home range, 232, 410, 489
- hunter recruitment, 370
- hunter retention, 370
- hunting, 338
- hunting management, 267
- hunting season, 322
- hybridization, 121

- I
 - incentives, 348
 - insecticide, 78
 - invertebrate biomass, 16, 66
 - invertebrate richness, 16, 66

- J
 - Johnson's rank method, 160

- K
 - kernel estimator, 489

- L
 - landowners, 348
 - landscape composition, 432
 - landscape structure, 432

- M
 - management, 11, 251, 456
 - management philosophies, 146
 - mesomammal abundance, 447
 - metapopulation, 135
 - microsatellites, 121
 - miolo, 282
 - minimum convex polygon, 489
 - Minnesota, 302
 - Mississippi, 220
 - modern quail management, 338
 - Montana, 260
 - Montezuma quail, 322, 426
 - morning covey call, 38
 - mortality, 54, 467

- N
 - Nothura maculosa*, 111
 - necklace-type collar transmitter, 480
 - nest, 256
 - nesting success, 447
 - New Mexico, 456
 - northern bobwhite, 16, 38, 46, 54, 66, 78, 88, 121, 135, 146, 160, 178, 184, 193, 220, 232, 251, 256, 271, 282, 315, 328, 338, 348, 396, 420, 432, 447, 467, 489, 518

- O
 - Ouachita Highlands, 146

- P
 - Perdix perdix*, 24, 27, 291, 380, 391
 - Phasianus colchicus*, 302, 410, 505
 - Procyon lotor*, 256
 - pairs, 172
 - parent fostering, 391
 - Partridge Count Scheme, 27
 - passive management, 370
 - pen-raised quail, 396
 - pen-reared, 121
 - pheasant, 260, 505
 - pine-bluestem, 184
 - pine-grassland restoration, 146
 - plantation, 338
 - population, 420
 - population dynamics, 267
 - population estimate, 99
 - population recovery, 291
 - pre-season release, 396
 - predation, 24, 256, 456
 - predation control, 24
 - predator control, 447
 - PREFER, 160
 - prescribed burning, 184, 271
 - prescribed fire, 78, 146
 - PROC NL MIXED, 232
 - productivity, 271
 - Program MARK, 232
 - proximal primary feather, 505

- Q
 - quail, 88, 178, 282, 370

- R
 - raccoon, 256
 - radio telemetry, 426
 - radio-handicapping, 518
 - radio-marked, 426
 - radio-telemetry, 338
 - radiotelemetry, 456
 - radius of audibility, 38
 - re-introduction, 391
 - red imported fire ant, 78
 - red-legged partridge, 267

reintroduction, 380
 releasing, 380
 reproduction, 232
 research, 370
 research and monitoring, 2
 restocking, 380
 ring-necked pheasant, 302
 riparian vegetation, 106
 Rolling Plains, 447
 ruffed grouse, 499
 ruffed grouse chicks, 480
 rural abandonment, 267

S

Solenopsis invicta, 78
Sorghum vulgare, 282
 salt cedar, 106
 sample size, 46
 SAS, 232
 scale, 135
 scale of management, 271
 scaled quail, 210, 456
 September burn, 315
 sex, 135
 simulation, 46
 sociological, 322
 South Carolina, 396
 south Texas, 328
 southeastern USA, 54, 467, 518
 southern African gamebirds, 11
 space use, 220
 spotted tinamou, 111
 suitability, 220
 supplemental feed, 282
 supplemental feeding, 210
 surveillance camera, 210
 survey, 38
 survival, 54, 111, 121, 172, 232, 271, 391, 426, 432, 456, 467, 518
 sweetgum, 315
 Switzerland, 391

T

Texas, 420
 telemetry, 54, 467, 518
 Tennessee, 121
 Texas, 78, 106, 251, 322, 447, 456
 threat, 11
 timber, 178
 TOBEC, 499
 total body electrical conductivity, 499

Trans-Pecos, 106, 322

U

UK, 24, 27, 291
 Upper Coastal Plain, 348
 urbanization, 370
 usable space, 146, 184
 utilization, 11

W

water developments, 210
 weather, 193
 West Virginia, 480
 wild pheasant, 410
 wildlife stand improvement, 184
 winter habitat, 432
 woody control, 315