
2000

Restocking Northern Bobwhites in East Texas: A Genetic Assessment

Michael A. Nedbal
Texas A&M University

Steven G. Evans
Texas A&M University

Rodney L. Honeycutt
Texas A&M University

R. Montague Whiting Jr.
Austin State University

Donald R. Dietz
Texas A&M University

Follow this and additional works at: <https://trace.tennessee.edu/nqsp>

Recommended Citation

Nedbal, Michael A.; Evans, Steven G.; Honeycutt, Rodney L.; Whiting, R. Montague Jr.; and Dietz, Donald R. (2000) "Restocking Northern Bobwhites in East Texas: A Genetic Assessment," *National Quail Symposium Proceedings*: Vol. 4 , Article 28.

<https://doi.org/10.7290/nqsp045rxr>

Available at: <https://trace.tennessee.edu/nqsp/vol4/iss1/28>

This article is brought to you freely and openly by Volunteer, Open-access, Library-hosted Journals (VOL Journals), published in partnership with The University of Tennessee (UT) University Libraries. This article has been accepted for inclusion in National Quail Symposium Proceedings by an authorized editor. For more information, please visit <https://trace.tennessee.edu/nqsp>.

RESTOCKING NORTHERN BOBWHITES IN EAST TEXAS: A GENETIC ASSESSMENT

Michael A. Nedbal

Department of Wildlife and Fisheries Sciences, Texas A&M University, College Station, TX 77843

Steven G. Evans

Department of Wildlife and Fisheries Sciences, Texas A&M University, College Station, TX 77843

Rodney L. Honeycutt

Department of Wildlife and Fisheries Sciences, Texas A&M University, College Station, TX 77843

R. Montague Whiting, Jr.

College of Forestry, Stephen F. Austin State University, Nacogdoches, TX 75962

Donald R. Dietz

Temple-Inland Forest Products Corporation, Lufkin, TX 75904

ABSTRACT

Habitat on a 610-ha study area in the Pineywoods Ecological Region of eastern Texas was enhanced for northern bobwhites (*Colinus virginianus*). In February and March 1990, 1991, and 1992, bobwhites from south Texas (*C. v. texanus*) and disjunct areas of east Texas (*C. v. mexicanus*) were captured, radio tagged, and relocated to the study area which had a small (<25 birds) resident population. Blood samples were collected from the birds relocated in 1991. Samples were also collected from birds in the resident population during 1991 and 1992; these birds were assumed to be offspring of the previous years' resident and relocated bobwhites. Restriction site variation of mitochondrial DNA (mtDNA) revealed geographic subdivision between the subspecies but not between resident and east Texas relocated birds. The observed frequency differences of mtDNA haplotypes were used to assess the relative reproductive success of the 2 subspecies. Among the birds examined for mtDNA variation, offspring produced on the study area during 1990 and 1991 were genetically more similar to the east Texas subspecies than to the south Texas subspecies. These results suggest that efforts to restock northern bobwhite should involve either local birds or birds from the same subspecies. Management implications of these findings are discussed.

Citation: Nedbal, M.A., S.G. Evans, R.L. Honeycutt, R.M. Whiting, Jr., and D.R. Dietz. 2000. Restocking northern bobwhites in East Texas: a genetic assessment. Page 115 in L.A. Brennan, W.E. Palmer, L.W. Burger, Jr., and T.L. Pruden (eds.). Quail IV: Proceedings of the Fourth National Quail Symposium. Tall Timbers Research Station, Tallahassee, FL.