

2000

Influence of Habitat Type and Prescribed Burning on Feral Swine Depredation of Artificial Quail Nests

Douglas R. Tolleson

Texas Agricultural Experiment Station

William E. Pinchak

Texas Agricultural Experiment Station

Dale Rollins

Texas Agricultural Experiment Station

R. James Ansley

Texas Agricultural Experiment Station

Follow this and additional works at: <https://trace.tennessee.edu/nqsp>

Recommended Citation

Tolleson, Douglas R.; Pinchak, William E.; Rollins, Dale; and Ansley, R. James (2000) "Influence of Habitat Type and Prescribed Burning on Feral Swine Depredation of Artificial Quail Nests," *National Quail Symposium Proceedings*: Vol. 4 , Article 26.

<https://doi.org/10.7290/nqsp04way1>

Available at: <https://trace.tennessee.edu/nqsp/vol4/iss1/26>

This article is brought to you freely and openly by Volunteer, Open-access, Library-hosted Journals (VOL Journals), published in partnership with The University of Tennessee (UT) University Libraries. This article has been accepted for inclusion in National Quail Symposium Proceedings by an authorized editor. For more information, please visit <https://trace.tennessee.edu/nqsp>.

INFLUENCE OF HABITAT TYPE AND PRESCRIBED BURNING ON FERAL SWINE DEPREDATION OF ARTIFICIAL QUAIL NESTS

Douglas R. Tolleson¹

Texas Agricultural Experiment Station, Vernon, TX 76385

William E. Pinchak

Texas Agricultural Experiment Station, Vernon, TX 76385

Dale Rollins

Texas Agricultural Extension Service, San Angelo, TX 76901

R. James Ansley

Texas Agricultural Experiment Station, Vernon, TX 76385

ABSTRACT

Feral swine (*Sus scrofa*) have increased both spatially and numerically in the Rolling Plains of northwest Texas. Resource managers have little information on the interrelationships between feral swine and native species such as the northern bobwhite (*Colinus virginianus*). Nest predation by feral swine could adversely affect bobwhite reproduction. Honey mesquite (*Prosopis glandulosa*) and redberry juniper (*Juniperus pinchotii*) are the dominant woody plants in northwestern Texas and prescribed burning is used to manage both species. Two experiments were conducted to determine the extent of feral swine depredation on artificial quail nests. In 1992–93, we monitored nest loss in mesquite and juniper habitats for six weeks. Total depredation reached 85 and 98% after three and six weeks respectively, during 1992, compared to 60 and 92% in 1993. Feral swine depredation of artificial bobwhite nests was 33% across years and was evenly distributed between vegetation types. During three-week periods in June and July 1994, depredation was greater ($P < 0.001$) in an unburned (90%) pasture than in unburned areas within a burned (32%) pasture. Decreased predator activity in the burned pasture was probably due to temporary prey displacement and less forage. Feral swine depredation of artificial bobwhite nests was 14% in 1994. Feral swine could potentially have negative impacts for northern bobwhites in mesquite and juniper habitats of northwest Texas.

Citation: Tolleson, D.R., W.E. Pinchak, D. Rollins, and R.J. Ansley. 2000. Influence of habitat type and prescribed burning on feral swine depredation of artificial quail nests. Page 110 in L.A. Brennan, W.E. Palmer, L.W. Burger, Jr., and T.L. Pruden (eds.). Quail IV: Proceedings of the Fourth National Quail Symposium. Tall Timbers Research Station, Tallahassee, FL.

¹ Present address: Texas A&M University System, Rangeland, Ecology and Management, Mail Stop 2126, College Station, TX 77843.